

Solution of the master equation for Bak-Sneppen model of biological evolution in finite ecosystem

Yu. M. Pis'mak

Department of Theoretical Physics

Saint-Petersburg State University

Ul'yanovskaya 1 Stary Peterhof 198904 Saint-Petersburg Russia

E-Mail pismak@phim.niif.spb.su

ABSTRACT

The master equations describing processes of biological evolution in the framework of the random neighbor Bak-Sneppen model are studied. For the ecosystem of N species they are solved exactly and asymptotical behavior of this solution for large N is analyzed.

1 Introduction

The model of biological evolution proposed by Bak and Sneppen [1,2] describes mutation and natural selection of interacting species. It is the dynamical system that is defined as follows. The state of the ecosystem of N species is characterized by a set $\{x_1, \dots, x_N\}$ of N number, $0 \leq x_i \leq 1$. In so doing, x_i represents the barrier toward further evolution of the species. Initially, each x_i is set to a randomly chosen value. At each time step the barrier x_i with minimal value and $K - 1$ other barriers are replaced by K new random numbers. In the random neighbor model (RNM), which will be considered in this paper the $K - 1$ replaced non-minimal barriers are chosen at random.

The RNM is the simplest model describing the avalanche-like processes, which are supposed by a conception of "punctuated equilibrium" in biological evolution. These processes are the most characterizing feature for self-organized criticality recently intensively investigated both numerically and analytically [3-6]. The RNM is more convenient for analytical studies. The master equations obtained in [3] for RNM are very useful for this aim. In [7] the explicit solution of master equations was found for infinite ecosystem. In this paper we solve the master equations for finite number N of species in ecosystem. We restrict ourselves with the simplest case $K = 2$.

2 Master equations for RNM

The master equations for the RNM are obtained in [3]. They are of the form:

$$P_n(t+1) = A_n P_n(t) + B_{n+1} P_{n+1}(t) + C_{n-1} P_{n-1}(t) + D_{n+2} P_{n+2}(t) + (B_1 \delta_{n,0} + A_1 \delta_{n,1} + C_1 \delta_{n,2}) P_0(t) \quad (1)$$

Here, $P_n(t)$ is the probability that at the time t is the number of barriers having values less than a fixed value λ at the time t ; $0 \leq n \leq N$, $0 \leq \lambda \leq N$, $0 \leq t$; $P_n(0)$

are proposed to be given. For $0 < n \leq N$

$$\begin{aligned} A_n &= 2\lambda(1 - \lambda) + \frac{n-1}{N-1}\lambda(3\lambda - 2), \\ B_n &= (1 - \lambda)^2 + \frac{n-1}{N-1}(1 - \lambda)(3\lambda - 1), \\ C_n &= \lambda^2 - \frac{n-1}{N-1}\lambda^2, \quad D_n = (1 - \lambda)^2 \frac{n-1}{N-1}, \end{aligned} \quad (2)$$

and $A_n = B_n = C_n = D_n = 0$ for $n = 0, n > N$.

In virtue of the definition of $P_n(t)$,

$$P_n(t) \geq 0, \quad (3)$$

$$\sum_{n=0}^N P_n(t) = 1. \quad (4)$$

Making summation in (1) over n and taking into account (2) it is easy to establish that

$$\sum_{n=0}^N P_n(t+1) = \sum_{n=0}^N P_n(t). \quad (5)$$

Therefore, if $P_n(0)$ are chosen in such a way that (4) is fulfilled for $t = 0$, then in virtue of (5) it is the case for the solution of (1) for $t > 0$ too. For analysis of (1) it is convenient to introduce the generating function $q(z, u)$:

$$q(z, u) \equiv \sum_{t=0}^{\infty} \sum_{n=0}^N P_n(t) z^n u^t. \quad (6)$$

In virtue of (3),(4) $q(z, t)$ is polynomial in z , analytical in u for $|u| < 1$ and

$$q(1, u) = \frac{1}{1 - u}. \quad (7)$$

The master equations (1) can be rewritten for the generating function $q(z, u)$ as follows:

$$\begin{aligned} \frac{1}{u}(q(z, u) - q(z, 0)) &= (1 - \lambda + \lambda z)^2 \left(\frac{1}{z} \left(1 - \frac{1 - z}{N - 1} \left(\frac{1}{z} - \frac{\partial}{\partial z} \right) \right) \times \right. \\ &\quad \left. \times (q(z, u) - q(0, u)) + q(0, u) \right). \end{aligned} \quad (8)$$

The function $q(z, 0) = \sum_{n=0}^N P_n(0) z^n$ in (8) is assumed to be given.

3 Asymptotic expansion of $q(z, u)$ for big N

If the function $q(z, 0)$ has an asymptotic expansion in the region of big N of the form:

$$q(z, 0) = \sum_{k=0}^{\infty} \frac{q_k(z, 0)}{(N-1)^k}$$

then the equation (8) enables one to obtain the similar asymptotic expansion for $q(z, u)$:

$$q(z, u) = \sum_{k=0}^{\infty} \frac{q_k(z, u)}{(N-1)^k}.$$

The main approximation of $q_0(z, u)$, the function $q(z, u)$ can be found from the equation

$$(z - u(1 - \lambda + \lambda z)^2)q_0(z, u) = zq_0(z, 0) + u(1 - \lambda + \lambda z)^2(z - 1)q_0(0, u) \quad (9)$$

following from (8). Since $q_0(z, u)$ is analytical for $|z| < 1, |u| < 1$,

$$0 = \alpha q_0(\alpha, 0) + u(1 - \lambda + \lambda \alpha)^2(\alpha - 1)q_0(0, u) \quad (10)$$

where

$$\alpha = \alpha(u) = \frac{1 - 2\lambda(1 - \lambda)u - (1 - 4\lambda(1 - \lambda u))^{\frac{1}{2}}}{2\lambda^2 u} \quad (11)$$

is the solution of equation $\alpha - u(1 - \lambda + \lambda \alpha)^2 = 0$. Obviously, for sufficient little $|u|$, $|\alpha| < 1$. Thus, from equation (10) the function $q_0(0, u)$ can be found:

$$q_0(0, u) = \frac{q_0(\alpha, 0)}{1 - \alpha}. \quad (12)$$

Substituting (12) in the right hand side of (9), one can find the solution in the following form:

$$q_0(z, u) = \frac{zq_0(z, 0)(1 - \alpha) + (z - 1)u(1 - \lambda + \lambda z)^2q_0(\alpha, 0)}{(z - u(1 - \lambda + \lambda z)^2)(1 - \alpha)}, \quad (13)$$

where $\alpha(u)$ is defined by (10).

For $k > 0$ the functions $q_k(z, u)$ are defined by recurrent relations

$$q_k(z, u) = \frac{u(1 - \lambda + \lambda z)^2(z - 1)q_k(\alpha, 0) + (1 - \alpha)zq_k(z, 0)}{(z - u(1 - \lambda + \lambda z)^2)(1 - \alpha)} + \frac{u(1 - \lambda + \lambda z)^2(r_{k-1}(z, u) + r_{k-1}(\alpha, u))}{z - u(1 - \lambda + \lambda z)^2}. \quad (14)$$

Here,

$$r_k(z, u) \equiv z \frac{\partial}{\partial z} \frac{q_k(z, u) - q_k(0, u)}{z}. \quad (15)$$

In virtue of (13), (14) the first correction to the lowest approximation (12) of $q(z, u)$ has the form

$$q_1(z, u) = \frac{u(1 - \lambda + \lambda z)^2(z - 1)q_1(\alpha, 0) + (1 - \alpha)zq_1(z, 0)}{(z - u(1 - \lambda + \lambda z)^2)(1 - \alpha)} + \frac{u(1 - \lambda + \lambda z)^2(r_0(z, u) + r_0(\alpha, u))}{z - u(1 - \lambda + \lambda z)^2}. \quad (16)$$

where

$$r_0(z, u) = z \frac{\partial}{\partial z} \frac{(1 - \alpha)q_0(z, u) + (u(1 - \lambda + \lambda z)^2 - 1)q_0(\alpha, 0)}{(z - u(1 - \lambda + \lambda z)^2)(1 - \alpha)}. \quad (17)$$

4 Exact form of $q(z, u)$

Let us introduce the quantity

$$Q(z, u) \equiv \frac{q(z, u) - q(0, u)}{z}. \quad (18)$$

It follows from (8) that this function fulfills the relation of the form:

$$\begin{aligned} (z - u(1 - \lambda + \lambda z)^2 - \frac{u(1 - \lambda + \lambda z)^2(1 - z)}{N - 1} \frac{\partial}{\partial z}) Q(z, u) = \\ = q(z, 0) + (u(1 - \lambda + \lambda z)^2 - 1)q(0, u). \end{aligned} \quad (19)$$

This inhomogeneous differential equation for $Q(z, u)$ has a special solution

$$Q(z, u) = (N - 1)e^{R(z, u)} \int_z^1 e^{-R(x, u)} g(x, u) dx \equiv Q_{sp}(z, u). \quad (20)$$

Here,

$$R(z, u) = \frac{N-1}{u} (\ln(1 - \lambda + \lambda z) - (1-u) \ln(1-z) + \frac{1-\lambda}{\lambda(1-\lambda+\lambda z)}), \quad (21)$$

$$g(x, u) = \frac{q(x, 0) + (u(1 - \lambda + \lambda z)^2 - 1)q(0, u)}{u(1 - \lambda + \lambda z)^2(1-x)} \quad (22)$$

and the derivative of $R(z, u)$ with respect to z has the form

$$\frac{\partial R(z, u)}{\partial z} = \frac{(N-1)(1 - (1 - \lambda + \lambda z)^2)}{u(1 - \lambda + \lambda z)^2(1-z)} \quad (23)$$

General solution of the corresponding to (19) homogeneous equation

$$(z - u(1 - \lambda + \lambda z)^2 - \frac{u(1 - \lambda + \lambda z)^2(1-z)}{N-1} \frac{\partial}{\partial z}) S(z, u) = 0 \quad (24)$$

is of the form

$$S(z, u) = F(u) e^{R(z, u)}. \quad (25)$$

Here, $F(u)$ is an arbitrary function of u . Hence, it follows from (19),(20),(25) that the function $Q(z, t)$ can be represented as follows:

$$Q(z, u) = F(u) e^{R(z, u)} + Q_{sp}(z, u). \quad (26)$$

In virtue of initial condition (7) for $q(z, u)$

$$Q(1, u) = \frac{1}{1-u} - q(0, u) \quad (27)$$

For $0 < u < 1, z \rightarrow 1$, $S(z, u)$ diverges and $S_{sp}(z, u)$ has the finite limit:

$$\lim_{z \rightarrow 1} Q_{sp}(z, u) = \frac{1}{1-u} - q(0, u) \quad (28)$$

Hence, $F(u) = 0$ in (25) and this representation for $Q(z, u)$ can be rewritten in the form:

$$\begin{aligned} Q(z, u) = (N-1) e^{R(z, u)} \int_{\frac{\lambda-1}{\lambda}}^1 e^{-R(x, u)} g(x, u) dx + \\ + (N-1) e^{R(z, u)} \int_z^{\frac{\lambda-1}{\lambda}} e^{-R(x, u)} g(x, u) dx \end{aligned} \quad (29)$$

It follows from (18) that

$$Q\left(\frac{\lambda-1}{\lambda}, u\right) = \frac{\lambda(q(\frac{\lambda-1}{\lambda}, u) - q(0, u))}{\lambda-1}$$

For the terms in the right hand side of (29) we have for $0 < u < 1$, $0 < \lambda < 1$:

$$\lim_{z \rightarrow \frac{\lambda-1}{\lambda} + 0} e^{R(z, u)} = +\infty, \quad (30)$$

$$\lim_{z \rightarrow \frac{\lambda-1}{\lambda} + 0} (N-1)e^{R(z, u)} \int_z^{\frac{\lambda-1}{\lambda}} e^{-R(x, u)} g(x, u) dx = \frac{\lambda(q(\frac{\lambda-1}{\lambda}, u) - q(0, u))}{\lambda-1}. \quad (31)$$

Therefore (29) can represent the function $Q(z, u)$ with necessary analytical properties only if

$$\int_{\frac{\lambda-1}{\lambda}}^1 e^{-R(x, u)} g(x, u) dx = 0 \quad (32)$$

This equation defines the function $q(0, u)$:

$$q(0, u) = \frac{\frac{\lambda-1}{\lambda} \int_{\frac{\lambda-1}{\lambda}}^1 e^{-R(x, u)} \frac{q(x, 0) dx}{(1-\lambda+\lambda x)^2(1-x)}}{\frac{\lambda-1}{\lambda} \int_{\frac{\lambda-1}{\lambda}}^1 e^{-R(x, u)} \frac{(1-u(1-\lambda+\lambda x)^2) dx}{(1-\lambda+\lambda x)^2(1-x)}} \quad (33)$$

Thus, we obtain from (29) the solution of equation (8) in the following form:

$$\begin{aligned} q(z, u) = & z \frac{N-1}{u} e^{R(z, u)} \int_z^{\frac{\lambda-1}{\lambda}} e^{-R(x, u)} \frac{q(x, 0) dx}{(1-\lambda+\lambda x)^2(1-x)} + \\ & + q(0, u) \left(1 + z \frac{N-1}{u} e^{R(z, u)} \int_z^{\frac{\lambda-1}{\lambda}} e^{-R(x, u)} \frac{(1-u(1-\lambda+\lambda x)^2) dx}{(1-\lambda+\lambda x)^2(1-x)}\right), \end{aligned} \quad (34)$$

where $q(0, u)$ is defined by (33).

5 Conclusion

We constructed the solution of the master equation (8) for the finite number N of species in the ecosystem. It can be proven that the main term (13) of its asymptotic

for large N coincides with the one obtained in [7]. Using (34) one can obtain all the known analytical results for RNM. One can hope that it helps to understand better the most important properties of the self-organized criticality processes.

References

- [1] P.Bak, K.Sneppen Phys. Rev. Lett. **71** (1993) 4083
- [2] H. Flyvbjerg, K.Sneppen, P.Bak Phys. Rev. Lett. **71** (1993) 4087
- [3] J. de Boer, B.Derrida, H. Flyvbjerg, A.D. Jacson, T.Wettig Phys. Rev. Lett. **73** (1994) 906
- [4] T.S. Ray, N.Jan Phys. Rev. Lett. **72** (1994) 4045
- [5] M.Paczuski, S.Maslov, P.Bak Europhys. Lett **27** (1994) 97
- [6] J. de Boer, A.D. Jackson, T.Wettig Criticality in Simple Models of Evolution, preprint cond-mat 9409074 (1994)
- [7] Yu.M. Pis'mak J. Phys. Math. Gen. **28** (1995) 3109